

A note on the action-angle variables for the rational Calogero–Moser system

Tomasz Brzeziński¹

Department of Mathematics, University of York
Heslington, York YO10 5DD, England

Cezary Gonera, Piotr Kosiński and Paweł Maślanka

Department of Theoretical Physics II, University of Łódź,
ul. Pomorska 149/153, 90-236 Łódź, Poland

Abstract

A relationship between the action-angle variables and the canonical transformation relating the rational Calogero–Moser system to the free one is discussed.

¹Lloyd's of London Tercentenary Fellow. On leave from: Department of Theoretical Physics, University of Łódź, ul. Pomorska 149/153, 90-236 Łódź, Poland. E-mail: tb10@york.ac.uk

The aim of this note is to answer the question of S. Ruijsenaars [1] concerning the relationship between the action-angle variables [2] for the rational Calogero-Moser model [3] and the equivalence of the latter to free particle systems described explicitly with the help of $sl(2, \mathbb{R})$ dynamical symmetry in [4].

We begin by recalling the construction of the canonical transformation in [4]. This construction is based on the observation that many features of the rational Calogero-Moser model with the Hamiltonian

$$H_{CM} = \sum_{i=1}^N \frac{p_i^2}{2} + \frac{g}{2} \sum_{i \neq j} \frac{1}{(q_i - q_j)^2},$$

where p_i, q_i are the canonical variables, and g is a coupling constant, can be explained in terms of the dynamical $sl(2, \mathbb{R})$ symmetry. Consider the following four functions on the phase space:

$$T_+ = \frac{1}{\omega} H_{CM}, \quad T_- \equiv \omega \sum_{i=1}^N \frac{q_i^2}{2}, \quad T_0 \equiv \frac{1}{2} \sum_{i=1}^N q_i p_i, \quad \tilde{T}_+ = \frac{1}{\omega} \sum_{i=1}^N \frac{p_i^2}{2},$$

where $\omega \neq 0$ is a parameter. One easily checks that each of the sets $\{T_+, T_-, T_0\}$ and $\{\tilde{T}_+, T_-, T_0\}$ spans the $sl(2, \mathbb{R})$ Lie algebra with respect to the Poisson brackets, i.e.,

$$\{T_0, T_{\pm}\} = \pm T_{\pm}, \quad \{T_-, T_+\} = 2T_0,$$

and

$$\{T_0, \tilde{T}_+\} = \tilde{T}_+, \quad \{T_0, T_-\} = -T_-, \quad \{T_-, \tilde{T}_+\} = 2T_0.$$

These $sl(2, \mathbb{R})$ algebras act on the phase space in the standard way by means of the Poisson brackets. The action can be integrated to the symplectic action of the $SL(2, \mathbb{R})$ group. In the construction of the transformation from the Calogero-Moser system to free particles an important role is played by the following one-parameter family of canonical transformations

$$\begin{aligned} q_k &\rightarrow e^{i\lambda T_1} * q_k \equiv \sum_{n=0}^{\infty} \frac{(i\lambda)^n}{n!} \{T_1, \dots \{T_1, q_k\} \dots\}, \\ p_k &\rightarrow e^{i\lambda T_1} * p_k \equiv \sum_{n=0}^{\infty} \frac{(i\lambda)^n}{n!} \{T_1, \dots \{T_1, p_k\} \dots\}, \end{aligned} \tag{1}$$

where $T_1 = \frac{i}{2}(T_+ + T_-)$. Since $T_1 = \frac{i}{2\omega}H_C$, where

$$H_C = \sum_{i=1}^N \left(\frac{p_i^2}{2} + \frac{\omega^2 q_i^2}{2} \right) + \frac{g}{2} \sum_{\substack{i,j=1 \\ i \neq j}}^N \frac{1}{(q_i - q_j)^2}$$

is the Hamiltonian of the Calogero model, the transformation (1) can be viewed as the time evolution generated by the Calogero Hamiltonian H_C , with the time $t = \lambda/2\omega$. On the other hand the transformation (1) is simply a rotation in the space spanned by $T_0, T_{\pm}$ about the axis T_1 by an angle λ . Thus for $\lambda = \pi$ (i.e., $t = \pi/2\omega$) we have $H_{CM} = \omega T_+ \rightarrow \omega e^{i\pi T_1} * T_+ = \omega T_-$. Next we can make a rotation in the space spanned by $T_0, T_-, \tilde{T}_+$ about the axis $\tilde{T}_1 = \frac{i}{2}(\tilde{T}_+ + T_-)$ through the angle $-\pi$. In particular, this will rotate T_- to $\tilde{T}_+$. Since the latter is proportional to the Hamiltonian of the free theory $H_0 = \sum_{i=1}^N \frac{p_i^2}{2}$, the canonical transformation obtained by the combination of two rotations transforms the rational Calogero-Moser model to the free particle theory, i.e.,

$$H_{CM} \rightarrow e^{-i\pi \tilde{T}_1} * (e^{i\pi T_1} * H_{CM}) = H_0.$$

Furthermore, this transformation sends the standard integrals of motion of the Calogero-Moser model $\frac{1}{m}\text{Tr}(L^m)$, $m = 1, \dots, N$, where L is the Lax matrix,

$$L_{jk} = \delta_{jk} p_k + (1 - \delta_{jk}) \frac{ig}{q_j - q_k}, \quad (2)$$

to their free counterparts (obtained by setting $g = 0$). The same applies to the functions $\text{Tr}(QL^m)$, $m = 1, \dots, N$, with $Q = \text{diag}(q_1, q_2, \dots, q_N)$.

The Ruijsenaars construction of the action-angle variables for the Calogero-Moser model can be most clearly explained in terms of the Hamiltonian reduction [5]. We now briefly recall how the reduction procedure can be applied to the Calogero-Moser model [6]. One starts with the space of pairs (A, B) of $N \times N$ hermitian matrices. This space is equipped with the symplectic form

$$\Omega = \text{Tr}(dB \wedge dA). \quad (3)$$

The action of the unitary group $U(N)$,

$$U \in U(N) : \quad (A, B) \rightarrow (UAU^\dagger, UBU^\dagger), \quad (4)$$

preserves the form Ω in (3) and thus is a symplectic action. The reduced phase space is obtained with the help of the momentum map equation

$$i[A, B] = g(I - \mathbf{v}^\dagger \otimes \mathbf{v}), \quad \mathbf{v} = (1, 1, \dots, 1).$$

Using the symplectic action of the group $U(N)$ in equation (4), one can fix a gauge in which $A = \text{diag}(q_1, q_2, \dots, q_N)$. In this gauge B is the Lax matrix in equation (2), and Ω takes the standard form $\Omega = \sum_{i=1}^N dp_i \wedge dq_i$. Thus we conclude that the Calogero-Moser model can be obtained by the Hamiltonian reduction of a simple dynamical system in Γ defined by the Hamiltonian $H = \frac{1}{2}\text{Tr}B^2$.


On the other hand the symplectic transformation [7] $A \rightarrow \tilde{A} = B$, $B \rightarrow \tilde{B} = -A$ preserves the momentum map. Following Ruijsenaars we can fix a gauge in which $\tilde{A} = B$ is diagonal, i.e., $\tilde{A} = \text{diag}(I_1, I_2, \dots, I_N)$. In this gauge the Hamiltonian $H = \frac{1}{2}\text{Tr}B^2$ is $H = \frac{1}{2}\sum_{i=1}^N I_i^2$. Clearly, the variables $I_1, \dots, I_N$ are constants of motion and together with the diagonal elements $-\phi_1, \dots, -\phi_N$ of $\tilde{B} = -A$ in this gauge, form the complete set of canonical variables. Thus we conclude that (ϕ_i, I_i) are the action-angle variables for the matrix model.

It is now not difficult to relate this construction of action-angle variables to that of the canonical map [4] recalled at the beginning of this note. The action of the $sl(2, \mathbb{R})$ symmetry on the reduced phase space can be lifted to Γ . Using the explicit form of the Poisson brackets induced by the canonical form Ω (3), $\{A_{ij}, B_{kl}\} = \delta_{il}\delta_{jk}$ one easily verifies that the functions

$$t_+ = \frac{1}{2\omega}\text{Tr}B^2, \quad t_- = \frac{\omega}{2}\text{Tr}A^2, \quad t_0 = \frac{1}{2}\text{Tr}AB,$$

generate the $sl(2, \mathbb{R})$ Lie algebra, i.e., $\{t_0, t_\pm\} = \pm t_\pm$, $\{t_-, t_+\} = 2t_0$. The relationship between the actions of $sl(2, \mathbb{R})$ on the unreduced and reduced phase spaces can be

summarised in the following commutative diagram:



Again using the explicit form of the Poisson brackets one finds that $t_0, t_{\pm}$ act linearly on A, B . This means that for any fixed k, l , (A_{kl}, B_{kl}) is an $sl(2, \mathbb{R})$ doublet. Therefore a general $sl(2, \mathbb{R})$ transformation of Γ can be represented as

$$\begin{pmatrix} A \\ \frac{1}{\omega}B \end{pmatrix} \rightarrow \begin{pmatrix} \alpha & \beta \\ \gamma & \delta \end{pmatrix} \begin{pmatrix} A \\ \frac{1}{\omega}B \end{pmatrix}, \quad \alpha\delta - \beta\gamma = 1. \quad (5)$$

The transformation (5) is a lift of the $sl(2, \mathbb{R})$ action on the reduced phase space. Thus, in particular, the lift of the canonical transformation induced by $e^{i\pi T_1}$ (cf. equation (1)), must be of the form (5). We have

$$\begin{pmatrix} A \\ \frac{1}{\omega}B \end{pmatrix} \rightarrow e^{i\pi t_1} * \begin{pmatrix} A \\ \frac{1}{\omega}B \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \begin{pmatrix} A \\ \frac{1}{\omega}B \end{pmatrix} = \begin{pmatrix} \frac{1}{\omega}B \\ -A \end{pmatrix}.$$

This shows that the Ruijsenaars procedure corresponds to the lifting of the construction of the canonical mapping of the Calogero-Moser system to free particles in [4]. One has to keep in mind, however, that the diagonal elements of B are viewed as momentum variables in the Ruijsenaars approach while in [4] they are proportional to the position variables. This explains the need for the additional transformation $e^{-i\pi \tilde{T}_1}$ which exchanges the momentum and position variables (and kills the factor ω).

The reasoning presented above explains also in a straightforward way why the functions $\text{Tr} L^n = \text{Tr} B^n$ and $\text{Tr}(QL^n) = \text{Tr}(AB^n)$ are transformed to their free counterparts, while it is no longer the case for $\text{Tr}(Q^m L^n)$, $m \geq 2$. The point is that the $\text{Tr} B^n$ and $\text{Tr}(AB^n)$ depend only on the eigenvalues of B and diagonal elements of A in the gauge in which B is diagonal, while the $\text{Tr}(Q^m L^n)$, $m \geq 2$ depend on non-diagonal elements of A too.

One can quantise the matrix theory on unreduced phase space Γ . Since the action of $sl(2, \mathbb{R})$ is linear, it can easily be implemented on the quantum level too. Then one can use the quantum Hamiltonian reduction [8] and carry the Ruijsenaars procedure over to the quantum case (for a different approach see [9]). At this point the main advantage of the procedure producing the symplectic map in [4] is that it can be immediately quantised.

Acknowledgments

This research is supported by the British Council grant WAR/992/147. The work of PK is supported by the grant KBN 2 P03B 134 16.

References

- [1] S.N.M. Ruijsenaars, private communication during the *Needs '99* conference.
- [2] S.N.M. Ruijsenaars, Commun. Math. Phys. **115** (1988), 127.
- [3] F. Calogero, J. Math. Phys. **12** (1971), 419;
F. Calogero, G. Marchioro, J. Math. Phys. **15** (1974), 1425;
J. Moser, Adv. Math. **16** (1975) 1.
- [4] T. Brzeziński, C. Gonera and P. Maślanka, Phys. Lett. A **254** (1999), 185.
- [5] D. Kazhdan, B. Kostant and S. Sternberg, Commun. Pure Appl. Math. **31** (1978), 481.
- [6] M.A. Olshanetsky and A.M. Perelomov, Phys. Rep. **71** (1981), 313; A. Gorsky, Theor. Math. Phys. **103** (1995), 681.
- [7] J. Moser [in:] *Dynamical Systems*, Progress in Mathematics No. 8, Birkhäuser, Boston, 1980, p. 233.
- [8] A.P. Polychronakos, Phys. Lett. B **266** (1991), 29;
A. Gorsky and N. Nekrasov, Nucl. Phys. B **414** (1994), 213; P. Etingof, J. Math. Phys. **36** (1995), 2636.

- [9] S.N.M. Ruijsenaars, Commun. Math. Phys. **110** (1987), 191.